

PRIOR AUTHORIZATION REQUEST Repatha

PATIENT:	Name	Prescriber:	Name			
			Address			
	City, State, Zip		City, State, Zip			
	D.O.B		Phone			
	Member ID:		Fax			
			NPI			
	Medication	on Requested: Qty R	equested:			
prescribed quantities of	a medication for can be provided.	your patient that requires Prior Authorization be Please complete the following questions then the	ests for coverage with the prescriber. You have before benefit coverage or coverage of additional fax this form to the toll free number listed below. It is determined based on the plan's rules.			
SEC	TION A: PI	ease answer the following questi	<u>ons</u>			
1.	□ Yes □ No	Is the requested medication being used of	concurrently with Praluent or Juxtapid?			
2.	□ Yes □ No	Is the requested medication being prescr	ibed by, or in consultation with, a			
		cardiologist; an endocrinologist; or a physical	sician who focuses in the treatment of			
		cardiovascular (CV) risk management an	d/or lipid disorders?			
3.	What is the indication or diagnosis?					
	Atherosclerotic cardiovascular disease (ASCVD) [clinical] – Please answer questions 16 – 26					
	Heterozygous familial hypercholesterolemia (HeFH) – Please answer questions 4 – 11 & 21 – 26					
	Homozygous familial hypercholesterolemia (HoFH) - Please answer questions 12 – 15 & 21 – 26 Primary Hyperlipidemia – Please answer questions 23 – 30					
	All other indications or diagnoses – Please specify					
4.			density lipoprotein cholesterol (LDL-C) level			
4.		greater than or equal to 190 mg/dL (that	is, prior to treatment with antihyperlipidemic			
_		agents)?	on of Hoffill by my totions in the law density			
5.	⊔ res ⊔ No		on of HeFH by mutations in the low-density			
			in B (APOB), proprotein convertase subtilisin			
		kexin type 9 (PCSK9) or low-density lipor	protein receptor adaptor protein 1			
		(LDLRAP1) gene?				
6.	□ Yes □ No	Has the patient been diagnosed with Hef Network criteria?	FH by the prescriber using the Dutch Lipid			
7.	☐ Yes ☐ No	Does the patient have a score of greater	than 5?			
8.	□ Yes □ No	Has the patient been diagnosed with Hel Broome criteria?	FH by the prescriber using the Simon			
9.	□ Yes □ No	Has the patient met the threshold for "de	finite" or "possible" familial			
		hypercholesterolemia?	·			
10.	□ Yes □ No	Does the patient have clinical manifestati	ons of HeFH?			

If you have any questions, call: 800-753-2851

PRIOR AUTHORIZATION REQUEST Repatha

11.	□ Yes □ No	Does the patient have a treated low-density lipoprotein cholesterol (LDL-C) level greater than or equal to 100 mg/dL (that is, after treatment with antihyperlipidemic agents but prior to treatment with PCSK9 inhibitor therapy such as Praluent or Repatha)?
12.	□ Yes □ No	• •
13.	□ Yes □ No	Does the patient have an untreated LDL-C level greater than 500 mg/dL (that is, prior to treatment with antihyperlipidemic agents)?
14.	□ Yes □ No	Does the patient have a treated LDL-C level of 300 mg/dL or greater (that is, after treatment with antihyperlipidemic agents but prior to agents such as Repatha, or Juxtapid)?
15.	☐ Yes ☐ No	Does the patient have patient has clinical manifestations of HoFH?
16.	□ Yes □ No	Has the patient had a previous myocardial infarction (MI) or history of an acute coronary syndrome (ACS)?
17.	☐ Yes ☐ No	Does the patient have a diagnosis of angina (stable or unstable)?
18.	☐ Yes ☐ No	Does the patient have a past history of stroke or transient ischemic attack (TIA)?
19.	☐ Yes ☐ No	Does the patient have peripheral arterial disease (PAD)?
20.	□ Yes □ No	Has the patient undergone a coronary or other arterial revascularization procedure in the past?
21.	□ Yes □ No	Has the patient tried one high-intensity statin therapy (that is, atorvastatin 40 mg or greater daily; rosuvastatin 20 mg or greater daily [as a single-entity or as a combination product]) for at least 8 weeks continuously?
22.	□ Yes □ No	Does the patient's low-density lipoprotein cholesterol (LDL-C) level after this treatment remain greater than or equal to 70 mg/dL?
23.	□ Yes □ No	Has the patient been determined to be statin intolerant by experiencing statin-related rhabdomyolysis?
24.	□ Yes □ No	Has the patient been determined to be statin intolerant by experiencing skeletal-related muscle symptoms?
25.	□ Yes □ No	Did the skeletal-related muscle symptoms occur while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products)?
26.	□ Yes □ No	When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) did the skeletal-related muscle symptoms resolve upon discontinuation of each respective statin therapy (atorvastatin AND rosuvastatin)?
27.	□ Yes □ No	equal to 300 Agatston units?
28.	□ Yes □ No	Has the patient tried one high-intensity statin therapy (that is, atorvastatin 40 mg or greater daily; rosuvastatin 20 mg or greater daily [as a single-entity or as a

If you have any questions, call: 800-753-2851



PRIOR AUTHORIZATION REQUEST Repatha

		combination product])?		
29.	□ Yes □ No	Was the high-intensity statin therapy (that is, atorvastatin 40 mg or greater daily; rosuvastatin 20 mg or greater daily [as a single-entity or as a combination product]) given with ezetimibe (as a single-entity or as a combination product) for at least 8 weeks continuously?		
30.	□ Yes □ No	Does the patient's low-density lipoprotein cholesterol (LDL-C) level after this treatment regimen remain greater than or equal to 100 mg/dL?		
Please document the diagnoses, symptoms, and/or any other information important to this review:				
SEC	TION B	Physician Signature		

PHYSICIAN SIGNATURE

DATE

FAX COMPLETED FORM TO: 877-251-5896

Disclaimer: An authorization is not a guarantee of payment. Member must be eligible at the time services are rendered. Services must be a covered Health Plan Benefit and medically necessary with prior

Disclaimer: An authorization is not a guarantee of payment. Member must be eligible at the time services are rendered. Services must be a covered Health Plan Benefit and medically necessary with prior authorization as per Plan policy and procedures.

Confidentiality: The information contained in this transmission is confidential and may be protected under the Health Insurance Portability and Accountability Act of 1996. If you are not the intended recipient any use, distribution, or copying is strictly prohibited. If you have received this facsimile in error, please notify us immediately and destroy this document.