

RX.PA.017.MPC Intravenous Immune Globulin (IVIG), Subcutaneous Immune Globulin (SCIG) & Intramuscular Immune Globulin (IMIG)

The purpose of this policy is to define the prior authorization process for all commercially-available, formulary IVIG and SCIG products.

DEFINITIONS

INCAT (Inflammatory Neuropathy Cause and Treatment Scale) – is used to assess functional disability of both upper and lower extremities in chronic inflammatory demyelinating polyradiculoneuropathy (CIDP).

Grade	Arm Disability
0	No upper limb problems
1	Symptoms, in one or both arms, not affecting the ability to perform any of the following functions: doing all zips and buttons; washing or brushing hair; using a knife and fork together; an handling small coins
2	Symptoms, in one arm or both arms, affecting but not preventing any of the above-mentioned functions
3	Symptoms, in one arm or both arms, preventing one or two of the above-mentioned functions
4	Symptoms, in one arm or both arms, preventing three or all of the functions listed, but some purposeful movements still possible
5	Inability to use either arm for any purposeful movement
Grade	Leg Disability
0	Walking not affected
1	Walking affected, but walks independently outdoors
2	Usually uses unilateral support (stick, single crutch, one arm) to walk outdoors
3	Usually uses bilateral support (stick, crutches, frame, two arms) to walk outdoors
4	Usually uses wheelchair to travel outdoors, but able to stand and walk a few steps with help
5	Restricted to wheelchair, unable to stand and walk a few steps with help

Inflammatory Rasch-built Overall Disability Scale (I-RODS) – intended to specifically assess activity and social participation limitations in patients with inflammatory neuropathies

Overall Disability Sum Score (ODSS) / Overall Neuropathy Limitations Scale (ONLS) – focuses on upper and lower limb functions and consists of a checklist for interviewing patients. The ODSS was the first scale designed to assess the limitations of patients with immune-mediated peripheral neuropathies. To reduce a possible ceiling effect, the ODSS was modified to include climbing stairs and running. The new measure is called the ONLS.

Grade	Arm Disability
0	Normal
1	Minor symptoms in one or both arms but not affecting any of the functions listed
2	Disability in one or both arms affecting but not preventing any of the functions listed
3	Disability in one or both arms preventing at least one but not all functions listed
4	Disability in both arms preventing all functions listed but purposeful movement still possible
5	Disability in both arms preventing all purposeful movements
Grade	Leg Disability
0	Walking/climbing stairs/running not affected
1	Walking/climbing stairs/running is affected, but gait does not look abnormal
2	Walks independently but gait looks abnormal
3	Requires unilateral support to walk 10 metres (stick, single crutch, one arm)
4	Requires bilateral support to walk 10 metres (sticks, crutches, crutch and arm, frame)
5	Requires wheelchair to travel 10 metres but able to stand and walk 1 metre with the help of one person
6	Restricted to wheelchair, unable to stand and walk 1 metre with the help of one person, but able to make some purposeful leg movements
7	Restricted to wheelchair or bed most of the day, unable to make any purposeful movements of the legs

Refractory Myasthenia Gravis Disease- unchanged or worse disease after corticosteroids and at least 2 other immunosuppressants, used in adequate doses for an adequate duration, with persistent symptoms or side effects that limit functioning, as defined by patient and physician.

The drugs, intravenous immune globulin (IVIG) and subcutaneous immune globulin (SCIG), are subject to the prior authorization process.

A. PROCEDURE

IVIG and SCIG are used to increase circulating levels of gamma globulin in certain immunoglobulin deficiency states and in treatment of a limited number of specified diseases.

FDA Approved Products and Indications (IVIG, SCIG, IMIG)

Product Name	Route of Administration	PID	ITP	CLL	CIDP	KD	MMN	DM	Hepatitis A	Measles	Varicella
Asceniv	IV	X									
Bivigam	IV	X									
Carimune NF	IV	X	X								
Cutaquig	SC	X									
Cuvitru	SC	X									
Flebogamma DIF	IV	X									
Gamastan S/D	IM								X	X	X
Gammagard Liquid	IV/SC	X					X				
Gammagard S/D	IV	X	X	X		X					
Gammaked	IV/SC	X	X		X						
Gammaplex	IV	X	X								
Gamunex-C	IV/SC	X	X		X						
Hizentra	SC	X			X						
Hyqvia	SC	X									
Octagam	IV	X	X					X			
Panzyga	IV	X	X		X						
Privigen	IV	X	X		X						
Xembify	SC	X									

FDA approved indications for specific Ig products:

- Primary Immunodeficiency Diseases (PID) [includes, but are not limited to, the humoral immune defect in common variable immunodeficiency (CVID), X-linked agammaglobulinemia, congenital agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies]
- Idiopathic thrombocytopenic purpura (ITP)
- Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)
- Kawasaki Disease (KD)

- Multifocal Motor Neuropath (MMN)
- Dermatomyositis (DM)

- IVIG products will not be approved for subcutaneous use unless FDA approved for that route of administration
- Requests for FDA approved indications should utilize the preferred associated specific Ig products listed in the table above

NOTE: Eviti reviews prior authorization requests for all oncology related indications for Ig products.

Must meet all of the criteria listed below:

FDA Approved Indications

1. For Primary Immunodeficiency

- Syndromes may include:
 - Common Variable Immunodeficiency (Hypogammaglobulinemia)
 - Congenital Agammaglobulinemia
 - Bruton's or X-linked Agammaglobulinemia
 - Severe Combined Immunodeficiency (SCID)
 - X-linked Hyper-IgM Syndrome
 - Wiskott-Aldrich Syndrome
 - Hypergammaglobulinemia Types
- Must be prescribed by or in consultation with an immunologist or hematologist
- Must have deficient antibody production, as evidenced through a documented IgG level $\leq 500\text{mg/dL}^*$
 - Requests with IgG levels $>500\text{mg/dL}$ require chart documentation that provides clinical rationale for the use of IVIG or SCIG (**NOTE:** Several primary immunodeficiencies do have normal levels of IgG with documented specific antibody deficiency)
 - *Note: Members under the age of 1 normally possess lower IgG levels
- Must have history of at least 1 bacterial infection directly attributable to this deficiency
- Approve for 1 year initially
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.

For Primary Immunodeficiency

- Syndromes may include:
 - Common Variable Immunodeficiency (Hypogammaglobulinemia)

2. For Idiopathic or Immune Thrombocytopenic Purpura (ITP)

(Platelet counts expressed per mm^3)

- Must be prescribed by a hematologist or oncologist
- For children with ITP:
 - Must have ONE of the following:
 - Active bleeding AND platelet count $<30,000$
 - Upcoming invasive surgery AND either platelet level below

- threshold designated for procedure (threshold must be provided with request) OR blood loss is expected
- Non-life-threatening mucosal bleeding and/or diminished quality of life AND documented previous inadequate response or intolerance to corticosteroids
- For adults with ITP:
 - Must have ONE of the following:
 - Active bleeding AND platelet count <30,000
 - Upcoming invasive surgery AND platelet level below threshold designated for procedure (threshold must be provided with request)
 - Platelet count <30,000 AND documented previous inadequate response or intolerance to corticosteroids
- For pregnant women with ITP
 - Must be pregnant and have ONE of the following:
 - Platelet count <50,000
 - Upcoming invasive surgery/procedure
 - History of splenectomy
 - Previously delivered infants with autoimmune thrombocytopenia
- Approve for 1 month
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.
- **DOSING NOTE:** Clinical guidelines recommend up to 1,000mg/kg/dose as a 1-time dose; dosage may be repeated if necessary.

3. For Kawasaki Disease

- Prescribed by or in consultation with a pediatric cardiologist or infectious disease physician
- Must be receiving aspirin concomitantly
- Must be requesting treatment within the first 10 days of illness
 - If greater than 10 days after illness onset, persistent signs of inflammation (e.g., persistent fever without explanation, elevated ESR or CRP, coronary artery aneurysms)
- Approve for 1 dose
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.
- **DOSING NOTE:** Clinical guidelines recommend 2 grams/kg

4. For HIV (Human Immunodeficiency Virus) in pediatric patients

- Must be prescribed by an immunologist or infectious disease specialist
- Must be <13 years old
- Must have CD4 count $\geq 200/\text{mm}^3$

- Must have ONE of the following
 - Recurrent (2 or more) serious bacterial infections such as bacteremia, meningitis, or pneumonia during a 1-year period despite administration of highly active antiretroviral therapy (HAART) and prophylactic sulfamethoxazole/trimethoprim (TMP-SMZ) or other antimicrobials
 - Hypogammaglobulinemia with an IgG <400mg/dL
 - Absence of detectable antibodies to common antigens, (measles, pneumococcal, and/or haemophilus influenzae Type B)
 - Bronchiectasis not optimally responsive to antibiotics and pulmonary therapy
 - A need for passive immunization for measles if Intramuscular Immune Globulin (IMIG) is contraindicated. IM injection contraindicated with severe thrombocytopenia or any coagulation disorder
- Approve for 1 year
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.
- **DOSING NOTE:** Clinical guidelines recommend 400mg/kg every 2-4 weeks

5. For Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)

- Must be prescribed by a neurologist
- Must have a diagnosis of CIDP
- Must provide documentation of electrodiagnostic testing (an EMG report)
- Must have moderate-to-severe functional disability
- Must provide a baseline disability score (using a validated disability scale, such as I-RODs, ODSS, ONLS, or INCAT)
- Approve for 3 months
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.

6. For Multifocal Motor Neuropathy

- Must be prescribed by a neurologist
- Must provide chart note documentation supporting a clinical examination of the member and BOTH of the following:
 - Progressive, asymmetric limb weakness over a course of at least 1 month
 - No objective sensory abnormalities except for minor vibration sense abnormalities in the lower limbs
- Must provide documentation of electrodiagnostic testing (an EMG report)
- Approve for 2 months
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.
- **DOSING NOTE:** Clinical guidelines recommend up to 2g/kg given over 2-5 days

7. For Dermatomyositis and Polymyositis (including juvenile)

- Must have dermatomyositis and polymyositis confirmed by biopsy

- Prescribed by or in consultation with a neurologist or rheumatologist
- Must have tried and failed or have a contraindication to both of the following:
 - Corticosteroids for 3 months
 - Concomitant adjuvant therapy for at least 3 months (azathioprine, methotrexate, cyclosporine)
- Approve for 2 months for either diagnosis
- Approval will be based off of FDA approved indication with specific associated IG products. Please refer to the table above.

8. GamaSTAN S/D

- Prophylaxis of Hepatitis A:
 - Must be prescribed by an infectious disease specialist
 - Must not present clinical manifestations of hepatitis A
 - Must not have been exposed to hepatitis A for more than 2 weeks
- Prophylaxis of Measles:
 - Must be prescribed by an infectious disease specialist
 - Must not have been exposed to measles for 6 days or more
 - Must not be given concurrently with the measles vaccine
- Prophylaxis of Varicella:
 - Must be prescribed by an infectious disease specialist
 - Must have been exposed to varicella within the last 10 days

Off-label Uses

1. For Guillain-Barre Syndrome

- Must start IVIG within 4 weeks of onset of neuropathic symptoms
- Must be unable to walk independently
- Prescribed by or in consultation with a neurologist
- Approve for 1 month
- May approve 1 additional month

2. For Systemic Lupus Erythematosus (SLE)

- Must have severe active SLE
- Must have tried and failed or have contraindications to ALL of the following:
 - Corticosteroids

- Antimalarials
- 1 additional immunosuppressant (azathioprine, cyclophosphamide, cyclosporine, methotrexate)
- Approve for 3 months

3. For Multiple Sclerosis (MS)

- Must be prescribed by a neurologist
- For acute exacerbations of MS:
 - Must have a trial and failure or have contraindications to corticosteroids and plasma exchange
 - Approve for 1 month
 - Additional authorizations for treatment are made on a case-by-case basis, are subject to the above criteria, and require chart documentation describing previous response to treatment and clinical rationale for re-treatment if IVIG did not provide a sufficient response.
- For chronic maintenance treatment of MS:
 - Must have relapsing, remitting type of MS
 - Must have a trial and failure (duration of at least 3 months) or have contraindications to ALL of the following:
 - At least one interferon [interferon beta-1a (Avonex®, Rebif®) or interferon beta- 1b (Betaseron®, Extavia®)]
 - Glatiramer (Copaxone®)
 - Fingolimod (Gilenya®)
 - No previous trials are required if:
 - Member is pregnant
 - Member is immunosuppressed or is having recurrent infections
 - Approve for 6 months

4. For Autoimmune Mucocutaneous Blistering Disease (AMBD)

- Must have one of the following supported by biopsy:
 - Pemphigus vulgaris
 - Pemphigus foliaceus
 - Bullous pemphigoid
 - Mucous membrane pemphigoid (a.k.a., cicatricial pemphigoid), OR
 - Epidermolysis bullosa acquisita
- Must have a trial and failure or have contraindications to corticosteroids and immunosuppressive agents
- EXCEPTION: In rapidly progressive, extensive, or debilitating cases IVIG may be approved along with corticosteroids or immunosuppressive agents
- Approve for 4 months

5. For Myasthenia Gravis Syndrome

- Must have a diagnosis of Myasthenia Gravis
- Must be prescribed by a neurologist
- For acute use:
 - Chart documentation of acute exacerbation and impaired function is required (e.g. respiratory insufficiency, inability to swallow)

- Approve for 1 month
- For temporary use as a bridge to immunotherapy:
 - Must have a history of myasthenia gravis exacerbation
 - Must be recently started (within 3 months) on immunosuppressant therapy (e.g. azathioprine, mycophenolate, cyclosporine, or tacrolimus)
 - Chart documentation of use as bridge therapy is required
 - Approve for 6 months
- For stabilization prior to surgery:
 - Must have a history of myasthenia gravis with current or previous difficulty with swallowing, speech, or respiratory involvement (e.g. shortness of breath or reduce force vital capacity on pre-op pulmonary function test). Chart documentation of symptoms is required.
 - IVIG infusion must be scheduled within 14 days of anticipated surgery date
 - Approve for 1 month
- For chronic use in refractory disease:
 - Must have an adequate trial with inadequate response, significant side effects/toxicity, or have a contraindication to both of the following:
 - Cholinesterase inhibitors- pyridostigmine or neostigmine
 - Corticosteroids
 - Must have an adequate trial of at least 3 months each with inadequate response, significant side effects/toxicity, or have a contraindication to TWO of the following:
 - Azathioprine
 - Mycophenolate mofetil
 - Cyclosporine
 - Tacrolimus
 - Approve for 6 months

6. For Parvovirus B19 Infection

- Must have documentation (e.g. Polymerase Chain Reaction test result) confirming presence of HPV-B19 infection
- Must have severe anemia defined as hemoglobin level <8ng/dL
- Must have low reticulocyte count defined as <8x10⁹/L
- Must have history of immunodeficiency due to suppressive medications or HIV
- Approve for 1 month
- **DOSING NOTE:** Clinical guidelines recommend 400-500mg for 5 days

7. For Renal and/or Pancreatic Transplant Desensitization in Combination with Rituxan

- Must be prescribed by a transplant specialist
- Must be age 18 or older
- Must be awaiting kidney and/or pancreas transplant requiring desensitization as defined by the following criteria:
 - For deceased donor transplants:
 - Panel reactive antibody (PRA) level >30% **OR**
 - PRA <30% with previous kidney and/or pancreas transplant
 - For living donor transplants:
 - Positive crossmatch **OR**

- Positive donor-specific antibody using Luminex® assay
- Approve for 1 course of treatment (2 doses)

8. For Renal Transplant Desensitization

- Must be prescribed by a transplant specialist
- Must be awaiting kidney transplant (either from a living or deceased donor) and requiring desensitization
- Approve for 4 months

9. For Renal Post-Transplant Rejection

- Must have received a renal transplant from a living donor with post-transplant rejection
- Approve for 1 month

10. For Allogenic or Hematopoietic Stem Cell Transplantation (HSCT) (or Bone Marrow Transplant)

- Must have severe hypogammaglobulinemia (IgG <400 mg/dL)
- Must have history of recurrent infections
- Approve for 6 months

11. For Autoimmune Hemolytic Anemia

- Must have warm-type diagnosis
- Must have a trial and failure or have contraindications to corticosteroids
- Approve for 1 month

12. For Stiff-Person Syndrome

- Must have a diagnosis of Stiff-Person Syndrome confirmed by electromyography (EMG) or elevated levels of glutamic acid decarboxylase (GAD)
- Must be prescribed by a neurologist
- Must have an adequate trial with inadequate responses, significant side effects/toxicity, or have contraindications to THREE of the following:
 - Corticosteroids
 - Antiepileptics
 - Benzodiazepines
 - Muscle relaxants
 - Gabapentin
- Approve for 4 months

13. Steven's Johnson Syndrome (SJS) or Toxic Epidermal Necrolysis (TEN)

- Initial approval is 1 month (2 doses)
- **DOSING NOTE:** Package insert/clinical guidelines recommend 1 g/kg/day in divided doses for 3- 4 days

14. CAR-T Therapy-Related Toxicity

- Must be prescribed by a hematologist, oncologist, or infectious disease specialist
- Must have Hypogammaglobinemia (IgG level <400mg/dL)
- Must have documentation of serious or recurrent infections (particularly bacterial)
- Initial approval is 1 year
- **DOSING NOTE:** Clinical guidelines recommend 400-500mg/kg monthly

B. Must be prescribed at a dose within the manufacturer's dosing guidelines (based on diagnosis, weight, etc) listed in the FDA approved labeling.

C. IV and subcutaneous immune globulin products will be considered investigational or experimental for any other use and coverage may be provided if it is determined that the use is a medically accepted indication supported by nationally recognized pharmacy compendia (AHFS-DI, DrugDex, Lexi-Drug, etc...) or at least two published peer-reviewed randomized controlled trials for the treatment of the diagnosis(es) for which it is prescribed. Abstracts (including meeting abstracts) are excluded from review consideration. These requests will be reviewed on a case by case basis to determine medical necessity.

Reauthorization Criteria:

All prior authorization renewals are reviewed to determine the Medical Necessity for continuation of therapy. Authorization may be extended based upon:

- MPC Renewal:
 - Chart documentation from the provider that the member's condition has stabilized or improved based upon the prescriber's assessment while on therapy and clinical rationale for re-treatment
 - FDA approved indications should use the specific associated IG products. Please refer to the table above.
 - No reauthorizations allowed for GamaSTAN S/D
- Renewal from Previous Insurer:
 - Members who have received prior approval (from insurer other than MPC), or have been receiving medication samples, should be considered under Initial Authorization Criteria.
 - Provider has a documented clinical response of the member's condition which has stabilized or improved based upon the prescriber's assessment.

Additional Reauthorization Criteria:

1. **Primary Immunodeficiency**
 - Reauthorization approval: 1 year
 - Must submit documentation from the prescriber of an updated IgG level (within 60 days of the renewal request)

2. **Idiopathic or Immune Thrombocytopenic Purpura (ITP)**

- Reauthorizations are granted on a case-by-case basis and are subject to the initial criteria
- Must provide documented initial response to IVIG therapy
- Documentation of continued thrombocytopenia <20,000 or < 30,000 and clinically significant bleeding
- **DOSING NOTE:** Clinical guidelines recommend up to 1,000mg/kg/dose
- Reauthorization approval: 3 months

3. **Kawasaki Disease**

- First reauthorization is granted (approval of only 1 dose) based upon documentation showing the member failed to respond to therapy
- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- Provider must submit documentation describing treatment response and clinical rationale for retreatment
- **DOSING NOTE:** Clinical guidelines recommend 2 grams/kg
- Reauthorization approval: 1 dose

4. **HIV (Human Immunodeficiency Virus) in pediatric patients**

- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- **DOSING NOTE:** Clinical guidelines recommend 400mg/kg every 2-4 weeks
- Reauthorization approval: 1 year

5. **Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)**

- Reauthorizations are granted based upon documentation from the prescriber indicating the member's condition has improved as a result of treatment as evidenced by improvement or stability in the member's baseline disability score (using a validated disability scale, such as I-RODS, ODSS, ONLS, or INCAT).
- Reauthorization approval: 1 year

6. **Multifocal Motor Neuropathy**

- **DOSING NOTE:** Clinical guidelines recommend up to 2g/kg given over 2-5 days
- Reauthorization approval: 1 year

7. **Dermatomyositis and Polymyositis (including juvenile)**

- Reauthorization approval for Dermatomyositis: 1 year
- Reauthorization approval for Polymyositis: 2 months

8. **Guillain-Barre Syndrome**

- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- Reauthorization approval: 1 month

9. **Systemic Lupus Erythematosus (SLE)**

- Reauthorization approval: 1 year

10. **Multiple Sclerosis (MS)**

- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- **For acute exacerbations of MS:**
 - Reauthorization approval: 1 month
- **For chronic maintenance treatment of MS:**
 - Reauthorization approval: 6 months

11. **Autoimmune Mucocutaneous Blistering Disease (AMBD)**

- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- Reauthorization approval: 4 months

12. **Myasthenia Gravis Syndrome**

- Reauthorizations are granted on a case by case basis and are subject to the initial criteria
- **For acute use:**
 - Reauthorization approval: 1 month
- **For temporary use as a bridge to immunotherapy:**
 - Reauthorization approval: 6 months
- **For stabilization prior to surgery:**
 - Reauthorization approval: 1 month
 - Must submit documentation on clinical rationale for reauthorization
- **For chronic use in refractory disease:**
 - Must have chart documentation describing previous response to treatment, including improvement in symptoms that limit daily function
 - Reauthorization approval: 6 months

13. **Parvovirus B19 Infection**

- Additional authorizations for treatment made on a case-by-case basis and are subject to the initial criteria
- **DOSING NOTE:** Clinical guidelines recommend 400-500mg for 5 days
- **Reauthorization approval:** 1 month
- **Maximum of 1 reauthorization per 365 days**

14. Renal and/or Pancreatic Transplant Desensitization in Combination with Rituxan

- Additional authorizations for treatment are subject to the initial criteria and are not granted until 6 months have passed since the initial treatment.
- Reauthorization approval: 1 course of treatment (2 doses)

15. Renal Transplant Desensitization

- Additional authorizations are not granted until 12 months have passed since the initial treatment.
- Reauthorization approval: 4 months

16. Renal Post-Transplant Rejection

- Reauthorization approval: 1 month

17. Allogenic or Hematopoietic Stem Cell Transplantation (HSCT) (or Bone Marrow Transplant)

- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- Reauthorization approval: 6 months

18. Autoimmune Hemolytic Anemia

- Additional authorizations for treatment are made on a case-by-case basis and are subject to the initial criteria
- Reauthorization approval: 1 month

19. Stiff-Person Syndrome

- Reauthorization for an additional 2 months of treatment may be made on a case-by-case basis and is subject to the initial criteria
- Reauthorization approval: 2 months
- Continued use beyond 6 months of therapy is not authorized.

20. Steven's Johnson Syndrome (SJS) or Toxic Epidermal Necrolysis (TEN)

- Reauthorizations are granted on a case-by-case basis and are subject to the initial criteria
- **DOSING NOTE:** Package insert/clinical guidelines recommend 1 g/kg/day in divided doses for 3- 4 days
- Documentation of a new episode
- Reauthorization approval: 1 month (2 doses)

21. CAR-T Therapy-Related Toxicity

- **DOSING NOTE:** Clinical guidelines recommend 400-500mg/kg monthly
- Reauthorization approval: 1 year

Limitations:

Length of Authorization (if above criteria met)	
Initial Authorization	Case-by-Case bases (see criteria above)
Reauthorization	Case-by-Case bases (see criteria above)

If the established criteria are not met, the request is referred to a Medical Director for review.

HCPCS Code(s):

Code	Description
J1459	Injection, immune globulin (Privigen), intravenous, nonlyophilized (e.g., liquid), 500 mg
J1556	Injection, immune globulin (Cuvitru), 100 mg
J1559	Injection, immune globulin (Hizentra), 100 mg
J1560	Injection, gamma globulin, intramuscular, over 10 cc
J1561	Injection, immune globulin, (Gamunex/Gamunex-C/Gammaked), nonlyophilized (e.g., liquid), 500 mg
J1566	Injection, immune globulin, intravenous, lyophilized (e.g., powder), not otherwise specified, 500 mg
J1568	Injection, immune globulin, (Octagam), intravenous, nonlyophilized (e.g., liquid), 500 mg
J1569	Injection, immune globulin, (Gammagard liquid), nonlyophilized, (e.g., liquid), 500 mg
J1572	Injection, immune globulin, (Flebogamma/Flebogamma Dif), intravenous, nonlyophilized (e.g., liquid), 500 mg
J1551	Injection, immune globulin (cutaquig), 100 mg
J1558	Injection, immune globulin (xembify), 100 mg
J1599	Injection, immune globulin, intravenous, non-lyophilized (e.g., liquid), not otherwise specified, 500 mg (Panzyga)
J1460	Injection, gamma globulin, intramuscular, 1 cc
J1560	Injection, gamma globulin, intramuscular, over 10 cc

REFERENCES

1. Immune globulin intravenous (human) In: Drug information for the health care professional. USPDI— Volume I. 25rd ed. Greenwood Village, CO: 2005.; Thomson Reuters Micromedex; 2012.
2. Knezevic-Maramica I, Kruskall MS. Intravenous immune globulins: an update for clinicians. *Transfusion*. 2003;43:1460-1480.
3. Carimune NF [package insert]. Kankakee, IL: ZLB Behring LLC (manufactured by ZLB Behring AG, Switzerland); January 2005.
4. Flebogamma 5% solution [package insert]. Los Angeles, CA: Grifols USA, Inc (manufactured by Instituto Grifols, SA, Barcelona, Spain); January 2004.
5. Gammagard Liquid [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; July 2012.
6. Gamunex-C [package insert]. Elkhart, IN: Bayer Corporation; October 2010.
7. Octagam [package insert]. Herndon, VA: Octapharma USA, Inc (manufactured by Octapharma Pharmazeutika, Vienna, Austria); March 2007. Gammagard® Liquid 10% [package insert]. Westlake Village, CA: Baxter Healthcare Corporation; December 2011. Buckley RH, Schiff RI. The use of intravenous immune globulin in immunodeficiency diseases. *NEJM*. 1991;325:110-117.
8. Neunert et al. ASH 2011 Evidence Based Practice Guideline for Idiopathic thrombocytopenia purpura. *Blood* April 2011. 4191-4204.
9. Cines DB, Blanchette VS. Immune thrombocytopenic purpura. *NEJM*. 2002 Mar 28;346:995-1008.
10. British Committee for Standards in Haematology General Haematology Task Force. Guidelines for the investigation and management of idiopathic thrombocytopenic purpura in adults, children and in pregnancy. *Br J Haematol*. 2003;120:574-596.
11. Vesely SK, Perdue JJ, Rizvi MA, et al. Management of adult patients with persistent idiopathic thrombocytopenic purpura following splenectomy: a systematic review. *Ann Intern Med*. 2004;140:112-120.
12. Buchanan GR, de Alarcon PA, Feig SA, et al. Acute idiopathic thrombocytopenic purpura – management in childhood. *Blood*. 1997;89:1464-1465.
13. Immune globulin (Drugdex® Drug Evaluation). Micromedex® Healthcare Series: Thomson Micromedex, Greenwood Village, Colorado. Accessed: January 2012
14. Newburger JW, Takahashi M, Gerber MA, et al; Committee on Rheumatic Fever, Endocarditis and Kawasaki Disease; Council on Cardiovascular Disease in the Young; American Heart Association; American Academy of Pediatrics. Diagnosis, treatment, and long-term management of Kawasaki disease: a statement for health professionals from the Committee on Rheumatic Fever, Endocarditis and Kawasaki Disease, Council on Cardiovascular Disease in the Young, American Heart Association. *Circulation*. 2004;110:2747-2771.
15. Ratko TA, Burnett DA, Foulke GE, et al. Recommendations for off-label use

- of intravenously administered immunoglobulin preparations. *JAMA*. 1995;273:1865-1870.
16. Ballow M. Intravenous immunoglobulins: clinical experience and viral safety. *J Am Pharm Assoc*. 2002;42:449-459.
 17. British Columbia Blood Coordinating Office. *IVIG utilization management handbook*. 1st Ed. British Columbia, Canada: Provincial Blood Coordinating Office; April 2002. <http://www.bloodlink.bc.ca/ivig.html> Accessed 03/14/2005.
 18. [No authors listed]. Intravenous immunoglobulin for the prevention of infection in chronic lymphocytic leukemia. A randomized, controlled clinical trial. Cooperative Group for the Study of Immunoglobulin in Chronic Lymphocytic Leukemia. *N Engl J Med*. 1988;319:902-907.
 19. Gamm H, Huber C, Chapel H, et al. Intravenous immune globulin in chronic lymphocytic leukaemia. *Clin Exp Immunol*. 1994;97 Suppl 1:17-20.
 20. Centers for Disease Control and Prevention; Infectious Disease Society of America; American Society of Blood and Marrow Transplantation. Guidelines for preventing opportunistic infections among hematopoietic stem cell transplant recipients. *MMWR Recomm Rep*. 2000 Oct;49(RR-10):1- 125, CE1-7.
 21. Sullivan KM, Storek J, Kopecky KJ, et al. A controlled trial of long-term administration of intravenous immunoglobulin to prevent late infection and chronic graft-vs.-host disease after marrow transplantation: clinical outcome and effect on subsequent immune recovery. *Biol Blood Marrow Transplant*. 1996;2:44-53.
 22. Mouthon L, Lortholary O. Intravenous immunoglobulins in infectious diseases: where do we stand? *Clin Microbiol Infect*. 2003;9:333-338.
 23. Centers for Disease Control and Prevention. Guidelines for preventing opportunistic infections among HIV-infected persons – 2002 recommendations of the U.S. Public Health Service and the Infectious Diseases Society of America. *MMWR*. 2002;51(No. RR-8):1-52.
 24. American Academy of Pediatrics. **Human Immunodeficiency Virus Infection**. In: Pickering LK, ed. *Red Book: 2003 Report of the Committee on Infectious Diseases*. 26th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2003:360-382.
 25. [No authors listed]. Antiretroviral therapy and medical management of pediatric HIV infection and 1997 USPHS/IDSA report on the prevention of opportunistic infections in persons infected with human immunodeficiency virus. *Pediatrics*. 1998;102(4 Pt 2):999-1085.
 26. Mofenson LM, Oleske J, Serchuck L, et al; CDC; National Institutes of Health; Infectious Diseases Society of America. Treating opportunistic infections among HIV-exposed and infected children: recommendations from CDC, the National Institutes of Health, and the Infectious Diseases Society of America. *MMWR Recomm Rep*. 2004 Dec 3;53(RR-14):1-92.
 27. Spector SA, Gelber RD, McGrath N, et al. A controlled trial of intravenous immune globulin for the prevention of serious bacterial infections in children receiving

- zidovudine for advanced human immunodeficiency virus infection. Pediatric AIDS Clinical Trials Group. *N Engl J Med*. 1994;331:1181- 1187.
28. Ahmed AR, Dahl MV. Consensus statement on the use of intravenous immunoglobulin therapy in the treatment of autoimmune mucocutaneous blistering diseases. *Arch Dermatol*. 2003;139:1051-1059.
 29. Ruetter A, Luger TA. Efficacy and safety of intravenous immunoglobulin for immune-mediated skin disease. Current view. *Am J Clin Dermatol*. 2004;5:153-160.
 30. Dalakas MC. Intravenous immunoglobulin in autoimmune neuromuscular diseases. *JAMA*. 2004;291:2367-2375.
 31. Hadden RD, Hughes RA. Management of inflammatory neuropathies. *J Neurol Neurosurg Psychiatry*. 2003;74 Suppl 2:ii9-ii14.
 32. Danieli MG, Cappelli M, Malcangi G, et al. Long term effectiveness of intravenous immunoglobulin in Churg-Strauss syndrome. *Ann Rheum Dis*. 2004;63:1649-1654.
 33. Tsurikisawa N, Taniguchi M, Saito H, et al. Treatment of Churg-Strauss syndrome with high-dose intravenous immunoglobulin. *Ann Allergy Asthma Immunol*. 2004;92:80-87.
 34. Ljungman P, Reusser P, de la Camara R, et al; for the Infectious Diseases Working Party of the European Group for Blood and Marrow Transplantation. Management of CMV infections: recommendations from the infectious diseases working party of the EBMT. *Bone Marrow Transplant*. 2004;33:1075-1081.
 35. Sokos DR, Berger M, Lazarus HM. Intravenous immunoglobulin: appropriate indications and uses in hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2002;8:117-130.
 36. John R, Lietz K, Burke E, et al. Intravenous immunoglobulin reduces anti-HLA alloreactivity and shortens waiting time to cardiac transplantation in highly sensitized left ventricular assist device recipients. *Circulation*. 1999;100[suppl II]:II-229-II235.
 37. Itescu S, Burke E, Lietz K, et al. Intravenous pulse administration of cyclophosphamide is an effective and safe treatment for sensitized cardiac allograft recipients. *Circulation*. 2002;105:1214-1219.
 38. Jordan SC, Tyan D, Stablein D, et al. Evaluation of intravenous immunoglobulin as an agent to lower allosensitization and improve transplantation in highly sensitized adult patients with end-stage renal disease: report of the NIH IG02 trial. *J Am Soc Nephrol*. 2004;15:3256-3262.
 39. Mackay MT, Weiss SK, Adams-Webber T, et al; American Academy of Neurology; Child Neurology Society. Practice parameter: medical treatment of infantile spasms: report of the American Academy of Neurology and the Child Neurology Society. *Neurology*. 2004;62:1668-1681.
 40. Dykes AC, Walker ID, Lowe GD, et al. Combined prednisolone and intravenous immunoglobulin treatment for acquired factor VIII inhibitors: a 2-year review. *Haemophilia*. 2001;7:160-163.

41. Collins PW. Management of acquired haemophilia A – more questions than answers. *Blood Coagul Fibrinolysis*. 2003;14 Suppl 1:S23-27.
42. Rubinger M, Rivard GE, Teitel J, et al, the Inhibitor Subcommittee of the Association of Hemophilia Clinic Directors of Canada (AHCDC). Suggestions for the management of hemophiliacs and non- hemophiliacs with factor VIII inhibitors. 3rd ed. May 14, 1999. <http://www.ahcdc.ca/Suggestions.htm> Accessed 3/10/2005.
43. Baschieri L, Antonelli A, Nardi S, et al. Intravenous immunoglobulin versus corticosteroid in treatment of Graves' ophthalmopathy. *Thyroid*. 1997;7:579-585.
44. Kahaly G, Pitz S, Muller-Forell W, et al. Randomized trial of intravenous immunoglobulins versus prednisolone in Graves' ophthalmopathy. *Clin Exp Immunol*. 1996;106:197-202.
45. Hughes RA, Wijdicks EF, Barohn R, et al; Quality Standards Subcommittee of the American Academy of Neurology. Practice parameter: immunotherapy for Guillain-Barre syndrome: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2003;61:736-740.
46. Jahnke L, Applebaum S, Sherman LA, et al. An evaluation of intravenous immunoglobulin in the treatment of human immunodeficiency virus-associated thrombocytopenia. *Transfusion*. 1994;34:759- 764.
47. Majluf-Cruz A, Luna-Castanos G, Huitron S, et al. Usefulness of a low-dose intravenous immunoglobulin regimen for the treatment of thrombocytopenia associated with AIDS. *Am J Hematol*. 1998;59:127-132.
48. WinRho SDF® [package insert]. Westlake Village, CA (manufactured by Cangene Corporation, Winnipeg, Canada): October 2004.
49. Comi G, Roveri L, Swan A, et al; Inflammatory Neuropathy Cause and Treatment Group. A randomised controlled trial of intravenous immunoglobulin in IgM paraprotein associated demyelinating neuropathy. *J Neurol*. 2002;249:1370-1377.
50. Cherin P, Pelletier S, Teixeira A, et al. Intravenous immunoglobulin for dysphagia of inclusion body myositis. *Neurology*. 2002;58:326.
51. Jordan SC, Vo AA, Nast CC, et al. Use of high-dose human intravenous immunoglobulin therapy in sensitized patients awaiting transplantation: the Cedars-Sinai experience. *Clin Transpl*. 2003;:193- 198.
52. Jordan S, Cunningham-Rundles C, McEwan R. Utility of intravenous immune globulin in kidney transplantation: efficacy, safety, and cost implications. *Am J Transplant*. 2003;3:653-664.
53. Maddison P, Newsom-Davis J. Treatment for Lambert-Eaton myasthenic syndrome. *Cochrane Database Syst Rev*. 2003;(2):CD003279.
54. Achiron A, Kishner I, Sarova-Pinhas I, et al. Intravenous immunoglobulin treatment following the first demyelinating event suggestive of multiple sclerosis: a randomized, double-blind, placebo-controlled

- trial. *Arch Neurol*. 2004;61:1515-1520.
55. Gray O, McDonnell GV, Forbes RB. Intravenous immunoglobulins for multiple sclerosis. *Cochrane Database Syst Rev*. 2003;(4):CD002936.
 56. Goodin DS, Frohman EM, Garmany GP Jr, et al; Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and the MS Council for Clinical Practice Guidelines. Disease modifying therapies in multiple sclerosis: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and the MS Council for Clinical Practice Guidelines. *Neurology*. 2002;58:169-178.
 57. Achiron A, Kishner I, Dolev M, et al. Effect of intravenous immunoglobulin treatment on pregnancy and postpartum-related relapses in multiple sclerosis. *J Neurol*. 2004;251:1133-1137.
 58. Bux J, Behrens G, Jaeger G, et al. Diagnosis and clinical course of autoimmune neutropenia in infancy: analysis of 240 cases. *Blood*. 1998;91:181-186.
 59. Herrod HG. Management of the patient with IgG subclass deficiency and/or selective antibody deficiency. *Ann Allergy*. 1993;70:3-8.
 60. Lawton AR. IgG subclass deficiency and the day-care generation. *Pediatr Infect Dis J*. 1999;18:462-466.
 61. Buckley RH. Immunoglobulin G subclass deficiency: fact or fancy? *Curr Allergy Asthma Reports*. 2002;2:356-360.
 62. Silk HJ, Ambrosino D, Geha RS. Effect of intravenous gammaglobulin therapy in IgG2 deficient and IgG2 sufficient children with recurrent infections and poor response to immunization with Hemophilus influenzae type b capsular polysaccharide antigen. *Ann Allergy*. 1990;64:21-25.
 63. Bjorkander J, Bengtsson U, Oxelius VA, et al. Symptoms in patients with lowered levels of IgG subclasses, with or without IgA deficiency, and effects of immunoglobulin prophylaxis. *Monogr Allergy*. 1986;20:157-163.
 64. Dalakas MC, Fujii M, Li M, et al. High-dose intravenous immune globulin for stiff-person syndrome. *N Engl J Med*. 2001;345:1870-1876.
 65. Ioannou Y, Isenberg DA. Current concepts for the management of systemic lupus erythematosus in adults: a therapeutic challenge. *Postgrad Med J*. 2002;78:599-606.
 66. O'Donnell BF, Barr RM, Kobza A, et al. Intravenous immunoglobulin in autoimmune chronic urticaria. *Br J Dermatol*. 1998;138:101-106.
 67. Asero R. Are IVIG for chronic unremitting urticaria effective? *Allergy*. 2000;55:1099-1101.
 68. Dawn G, Urcelay M, Ah-Weng A, et al. Effect of high-dose intravenous immunoglobulin in delayed pressure urticaria. *Br J Dermatol*. 2003;149:836-840.
 69. Joint Tasks Force on Practice Parameters. The diagnosis and management of urticaria: a practice parameter part I: acute urticaria/angioedema part II: chronic urticaria/angioedema. *Ann Allergy Asthma Immunol* 2000;85(6 Pt 2):521-544.

70. Becker MD, Rosenbaum JT. Current and future trends in the use of immunosuppressive agents in patients with uveitis. *Curr Opin Ophthalmol*. 2000;11:472-477.
71. LeHoang P, Cassoux N, George F, et al. Intravenous immunoglobulin (IVIg) for the treatment of birdshot retinochoroidopathy. *Ocul Immunol Inflamm*. 2000;8:49-57.
72. Jayne DR, Chapel H, Adu D, et al. Intravenous immunoglobulin for ANCA-associated systemic vasculitis with persistent disease activity. *QJM*. 2000;93:433-439.
73. Relkin N, Szabo, Adamiak B, et al. Intravenous immunoglobulin (IVIG) treatment causes dose- dependent alterations in β -amyloid (A β) levels and anti-A β antibody titers in plasma and cerebrospinal fluid (CSF) of Alzheimer's disease (AD) patients [abstract]. Presented at: American Academy of Neurology 57th Annual Meeting; April 12, 2005: Miami Beach, FL.
74. Dodel RC, Du Y, Depboylu C, et al. Intravenous immunoglobulins containing antibodies against beta- amyloid for the treatment of Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 2004;75:1472-1474.
75. Hack CE, Scheltens P. Intravenous immunoglobulins: a treatment for Alzheimer's disease? *J Neurol Neurosurg Psychiatry*. 2004;75:1374-1375.
76. Niven AS, Argyros G. Alternate treatments in asthma. *Chest*. 2003;123:1254-1265.
77. Jolles S, Sewell C, Webster D, et al. Adjunctive high-dose intravenous immunoglobulin treatment for resistant atopic dermatitis: efficacy and effects on intracellular cytokine levels and CD4 counts. *Acta Derm Venereol*. 2003;83:433-437.
78. Paul C, Lahfa M, Bachelez H, et al. A randomized controlled evaluator-blinded trial of intravenous immunoglobulin in adults with severe atopic dermatitis. *Br J Dermatol*. 2002;147:518-522.
79. Jolles S. A review of high-dose intravenous immunoglobulin treatment for atopic dermatitis. *Clin Exp Dermatol*. 2002;27:3-7.
80. Jolles S, Hughes J. Importance of trial design in studies using high-dose intravenous immunoglobulin. *Br J Dermatol*. 2003;148:1284-5; author reply 1285-6.
81. Wakim M, Alazard M, Yajima A, et al. High dose intravenous immunoglobulin in atopic dermatitis and hyper-IgE syndrome. *Ann Allergy Asthma Immunol*. 1998;81:153-158.
82. Noh G, Lee KY. Intravenous immune globulin (i.v.IG) therapy in steroid-resistant atopic dermatitis. *J Korean Med Sci*. 1999;14:63-68.
83. Hanifin JM, Cooper KD, Ho V, et al. Guidelines of care for atopic dermatitis. *J Am Acad Dermatol*. 2004;50:391-404.
84. Wolff SN, Fay JW, Herzig RH, et al. High-dose weekly intravenous immunoglobulin to prevent infections in patients undergoing autologous bone marrow transplantation or severe myelosuppressive therapy. A study of the American Bone Marrow Transplant Group. *Ann Intern Med*. 1993;118:937-942.
85. Seider N, Beiran I, Scharf J, et al. Intravenous immunoglobulin therapy for resistant ocular Behcet's disease. *Br J Ophthalmol*. 2001;85:1287-1288.

86. Cordonnier C, Chevret S, Legrand M, et al; GREFIG Study Group. Should immunoglobulin therapy be used in allogeneic stem-cell transplantation? A randomized, double-blind, dose effect, placebo- controlled, multicenter trial. *Ann Intern Med.* 2003;139:8-18.
87. Gottstein R, Cooke RW. Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn. *Arch Dis Child Fetal Neonatal Ed.* 2003;88:F6-10.
88. Hommes OR, Sorensen PS, Fazekas F, et al. Intravenous immunoglobulin in secondary progressive multiple sclerosis: randomised placebo-controlled trial. *Lancet.* 2004;364:1149-1156.
89. Ohlsson A, Lacy JB. Intravenous immunoglobulin for suspected or subsequently proven infection in neonates. *Cochrane Database Syst Rev.* 2004:CD001239.
90. Ohlsson A, Lacy JB. Intravenous immunoglobulin for preventing infection in preterm and/or low-birth- weight infants. *Cochrane Database Syst Rev.* 2004;(1):CD000361.
91. Fanaroff AA, Korones SB, Wright LL, et al. A controlled trial of intravenous immune globulin to reduce nosocomial infections in very-low-birth-weight infants. National Institute of Child Health and Human Development Neonatal Research Network. *N Engl J Med.* 1994;330:1107-1113.
92. American College of Obstetricians and Gynecologists (ACOG). ACOG Practice Bulletin. Management of recurrent early pregnancy loss. Washington, DC: ACOG; 2001 Feb.
93. Branch DW, Peaceman AM, Druzin M, et al. A multicenter, placebo-controlled pilot study of intravenous immune globulin treatment of antiphospholipid syndrome during pregnancy. The Pregnancy Loss Study Group. *Am J Obstet Gynecol.* 2000;182(1 Pt 1):122-127.
94. Practice Committee of the American Society for Reproductive Medicine. Intravenous immunoglobulin (IVIG) and recurrent spontaneous pregnancy loss. *Fertil Steril.* 2004;82 Suppl 1:S199-200.
95. Levy Y, Amital H, Langevitz P, et al. Intravenous immunoglobulin modulates cutaneous involvement and reduces skin fibrosis in systemic sclerosis: an open-label study. *Arthritis Rheum.* 2004;50:1005- 1007.
96. White B, Bauer EA, Goldsmith LA, et al. Guidelines for clinical trials in systemic sclerosis (scleroderma). I. Disease-modifying interventions. The American College of Rheumatology Committee on Design and Outcomes in Clinical Trials in Systemic Sclerosis. *Arthritis Rheum.* 1995;38:351-360.
97. Pereyra F, Rubin RH. Prevention and treatment of cytomegalovirus infection in solid organ transplant recipients. *Curr Opin Infect Dis.* 2004;17:357-361.

98. Preiksaitis JK, Brennan DC, Fishman J, et al. Canadian Society of Transplantation consensus workshop on cytomegalovirus management in solid organ transplantation final report. *Am J Transplant.* 2005;5:218-227.
99. Hizentra™ [package insert]. Bern, Switzerland: CSL Behring; March 2010.
100. Gammaplex® [package insert]. Elstree Hertfordshire, United Kingdom: Bio Products Laboratory; 2009.
101. Shehata N, et al. The use of immunoglobulin therapy for patients with Primary Immune Deficiency: an evidence-based practice guideline. *Transfusion Medicine Reviews* 2010. 24(1): S28- S50. Yellen ES, et al. Performance of 2004 American Heart Association recommendations for treatment of Kawasaki disease. *Pediatrics* 2010. 125(2):e234-e241.
102. Mofenson L, et al. Guidelines for the Prevention and Treatment of Opportunistic Infections Among HIV-Exposed and HIV-Infected Children: Recommendations from CDC, the National Institutes of Health, the HIV Medicine Association of the Infectious Diseases Society of America, the Pediatric Infectious Diseases Society, and the American Academy of Pediatrics. *MMWR Recomm Rep.* 2009;10;58(RR-4):1-207.
103. Antoine JC, the French CIDP Study Group. Recommendations on diagnostic strategies for CIDP. *Journal of Neurology, Neurosurgery, Psychiatry* 2008. 79: 118.
104. Van Den Bergh PYK, et al. European Federation of Neurological Societies/PNS Guidelines on management of chronic inflammatory demyelinating polyradiculoneuropathy - first revision. *European Journal of Neurology* 2010. 17: 356-363.
105. Bromberg MB. Review of the Evolution of Electrodiagnostic Criteria for Chronic Inflammatory Demyelinating Polyradiculoneuropathy. *Issues & Opinions in Muscle & Nerve*; June 2011: 780-792.
106. Hughes RA, et al. Practice parameter: immunotherapy for Guillain-Barre syndrome: report of the Quality Standards Committee of the American Academy of Neurology. August 2, 2008.
107. Winer JB. When the Guillain Barre patient fails to respond to treatment. *Practice Neurology* 2009. 9:227-230.
108. Marie I, Mouthon L. Therapy of Polymyositis and dermatomyositis. *Autoimmunity Reviews* 2011. 11: 6-13.
109. Hak AE, et al. Dermatomyositis and polymyositis: new treatment targets on the horizon. *Journal of Medicine* 2011. 69(10): 415-416.
110. Aggarwal R, Oddis CN. Therapeutic Approaches in myositis. *Current Rheumatology Rep* 2011. 13: 182-191.
111. Berger JR. Functional improvement and symptom management in multiple sclerosis: clinical efficacy of current therapies. *American Journal of Managed Care* 2011. 17: S146-153.
112. Elovaara I, et al. Intravenous Immunoglobulins are a therapeutic option in the treatment of multiple sclerosis relapse. *Clinical Neuropharmacology* 2011. 34(2):84-88.
113. Rio J, et al. Multiple Sclerosis: current treatment algorithms. *Current Opinion in*

- Neurology 2011. 24: 230-237.
114. Feasby T, et al. Guidelines for the use of IVIG for neurological conditions. *Transfus Med Rev* 2007. 21 (2 suppl 1): S57-107.
 115. Gold R, et al. Drug Insight: the use of IVIG in neurology. *Nature Clinical Practice Neurology* 2007. 3:36-44.
 116. McGhee SA, et al. Persistent parvovirus-associated chronic fatigue treated with high dose IVIG. *Pediatric Infectious Disease Journal* 2005. 24(3):272-274.
 117. Baden LR, et al. NCCN: Prevention and Treatment of Cancer-Related Infections. Version 2.2011.
 118. Tomblyn M, et al. Guidelines for Preventing Infectious Complications among Hematopoietic Cell Transplantation Recipients. *Biol Blood Marrow Transplant* 2009. 15:1143-1238.
 119. Yildirim-Toruner C, Diamond B. Current and novel therapeutics in the treatment of systemic lupus erythematosus. *Clinical reviews in allergy and immunology* 2010. 303-312.
 120. Bayry J et al. Intravenous immunoglobulin treatment in rheumatic disease. *Nature Reviews Rheumatology* 2011. 7: 351-352.
 121. Petz LP. Treatment of autoimmune hemolytic anemia. *Current Opinion in Hematology*. 2001;8:411-416.
 122. Skele GO, et al. Guidelines for the treatment of autoimmune neuromuscular transmission disorders. *Euro J of Neurology* 2006. 13: 691-699.
 123. Fazekas et al. Randomised placebo-controlled trial of monthly intravenous immunoglobulin therapy in relapsing-remitting multiple sclerosis. *Lancet* 1997; 349:589-593.
 124. Achiron et al. Intravenous immunoglobulin treatment in multiple sclerosis: effect on relapses. *Neurology* 1998; 50: 398-402.
 125. Sorenson et al. Intravenous immunoglobulin G reduced MRI activity in relapsing multiple sclerosis. *Neurology* 1998 50: 1273-1281.
 126. Achiron et al. Intravenous immunoglobulin treatment following the first demyelinating event suggestive of Multiple Sclerosis. *Arch Neurology* 2004; 61:1515-1520
 127. Orange J, Hossny E, Weiler C, et al. Use of intravenous immunoglobulin in human disease: a review of evidence by members of the Primary Immunodeficiency Committee of the American Academy of Allergy, Asthma, and Immunology. *J Allergy Clin Immunol*. 2006;117(4):S525-S553.
 128. Leong H, Stachnik J, Bonk ME, Matuszewski KA. Unlabeled uses of intravenous immune globulin. *Am J Health Syst Pharm*. 2008;65(19)1815-1824.
 129. RhoGAM® and MICRhoGAM® [package insert]. Raritan, NJ; Ortho-Clinical Diagnostics, Inc.; 2007. Available at: <http://www.rhogam.com/Patient/News/Pages/default.aspx>.
 130. Panglobulin® [package insert]. Berne, Switzerland; ZLB Bioplasma AG; 2000. Available at: http://www.ctint.org/inserts/Panglobulin_insert.pdf.

131. Polygam® [package insert]. Westlake Village, CA; Baxter Healthcare Corporation; 2002. Available at: http://www.ctint.org/inserts/PolygamSD_insert.pdf.
132. Rhophylac® [package insert]. Bern, Switzerland; CSL Behring AG; 2010. Available at: <http://labeling.cslbehring.com/PI/US/Rhophylac/EN/Rhophylac-Prescribing-Information.pdf>.
133. Thymoglobulin® [package insert]. Cambridge, MA; Genzyme Corporation; 2008. Available at: http://www.thymoglobulin.com/home/thymo_pdf_pi.pdf.
134. Winrho® [package insert]. Winnipeg, Manitoba; Cangene Corporation; 2010. Available at: <http://www.winrho.com/pi.pdf>.
135. Patwa HS, Chaudry V, Katzberg H, et al. Evidence-based guideline: Intravenous immunoglobulin in the treatment of neuromuscular disorders: Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 2012;78:1009-1015
136. Leger J, Chassandre B, Musset L, et al. Intravenous immunoglobulin therapy in multifocal motor neuropathy: a double-blind, placebo-controlled study. *Brain* 2001;124:145-153
137. Federico P, Zochodne DW, Hahn AF, et al. Multifocal motor neuropathy improved by IVIg: randomized, double-blind, placebo-controlled study. *Neurology* 2000;55:1256-1262
138. Hughes R, Bensa S, Willison H, et al. Randomized controlled trial of intravenous immunoglobulin versus oral prednisolone in chronic inflammatory demyelinating polyradiculoneuropathy. *Ann Neurol* 2001;50:195-201
139. Hughes RA, Donofrio P, Bril V, et al. Intravenous immune globulin (10% caprylate- chromatography purified) for the treatment of chronic inflammatory demyelinating polyradiculoneuropathy (ICE study): a randomised placebo-controlled trial. *Lancet Neurol* 2008;136- 44
140. Mouthon L, Guillevin L, Tellier Z. Intravenous immunoglobulins in autoimmune- or parvovirus B19-mediated pure red-cell aplasia. *Autoimmunity Reviews* 2005;4:264-269
141. Crabol R, Terrier B, Rozenberg F, et al. Intravenous immunoglobulin therapy for pure red cell aplasia related to human parvovirus B19 infection: a retrospective study of 10 patients and review of the literature. *Clinical Infectious Disease* 2013;56(7):968-77
142. Umapathi T, Hughes RAC, Nobile-Orazio E, et al. Immunosuppressant and immunomodulatory treatments for multifocal motor neuropathy. *Cochrane Database of Systematic Reviews* 2012, Issue 4. Art. No.: CD003217. DOI: 10.1002/14651858.CD003217.pub4.

143. Karlson EW, Sudarsky L, Ruderman E, et al. Treatment of stiff-man syndrome with intravenous immune globulin. *Arthritis Rheum* 1994; 37:915.
144. Barker RA, Marsden CD. Successful treatment of stiff man syndrome with intravenous immunoglobulin. *J Neurol Neurosurg Psychiatry* 1997; 62:426.
145. Khanlou H, Eiger G. Long-term remission of refractory stiff-man syndrome after treatment with intravenous immunoglobulin. *Mayo Clin Proc* 1999; 74:1231.
146. Dalkas MC, Fujii M, Li M, et al. High-dose intravenous immune globulin for stiff-person syndrome. *N Engl J Med* 2001; 345:1870
147. International consensus guidance for management of myasthenia gravis June 2016 American Academy of Neurology
149. Neunert C, Terrell D, Arnold DM, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia. *Blood Advances*. 2019 Oct; 3(23): 3829-3866
150. McCrindle BW, Rowley AH, Newburger JW. Diagnosis, Treatment and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association. *Circulation*. 2017; 135:927-999. doi: 10.1161/CIR.0000000000000484
151. Panel on Opportunistic Infections in HIV-Exposed and HIV-Infected Children. Guidelines for the Prevention and Treatment of Opportunistic Infections in HIV-Exposed and HIV-Infected Children. Department of Health and Human Services. Available at https://clinicalinfo.hiv.gov/sites/default/files/inline-files/oi_guidelines_pediatrics.pdf. Last updated Nov 2013.
152. Joint Task Force of the EFNS and the PNS. European Federation of Neurological Societies/Peripheral Nerve Society Guideline on management of multifocal motor neuropathy. Report of a Joint Task Force of the European Federation of Neurological Societies and the Peripheral Nerve Society – first revision. *Journal of the Peripheral Nervous System* 2010;15:295-301.
151. Narayanaswami P, Sanders DB, Wolfe G, et al. International Consensus Guidance for Management of Myasthenia Gravis. *Neurology*® 2021;96:114-122. doi:10.1212/WNL.00000000000011124
154. Baden LR, et al. NCCN: Prevention and Treatment of Cancer-Related Infections. Version 1.2021.
155. Thompson, et al. NCCN: Management of Immunotherapy-Related Toxicities. Version 3.2021.
156. Poston JN, Gernsheimer TB. Management of Immune Thrombocytopenia in Pregnancy. *Annals of Blood*. 2021:

REVIEW HISTORY

DESCRIPTION OF REVIEW / REVISION	DATE APPROVED
<i>Annual Review</i>	02/2026
<i>Annual Review</i>	02/2025
<i>Annual Review</i> <i>Change in Non-MPC renewal to renewal from previous insurer</i>	02/2024
<i>Selected Revision: Update to reauthorization criteria based on case by case indications</i>	03/2023
<i>Annual Review</i>	02/2023
<i>Selected Revision: Update to FDA approved products per indication</i> <i>Addition of reauthorization criteria</i> <i>Addition of FDA approved indications for GamaSTAN S/D</i>	01/2023
<i>Update to off-label restrictions</i>	04/2022
<i>Annual review</i>	02/2022
<i>Addition of new indications for SJS/TEN and Car-T Therapy-Related Toxicity criteria update for: AMBD, CIDP, Guillain Barre, ITP, MN, Myasthenia Gravis Syndrome</i>	02/2022
<i>Addition of dosing requirements and off-label restrictions</i>	12/2021
<i>P&T Review</i>	11/2020