

RX.PA.067.MPC Ryplazim (plasminogen, human-tvmh)

The purpose of this policy is to define the prior authorization process for Ryplazim® (plasminogen, human-tvmh)

Ryplazim® (plasminogen, human-tvmh) is a plasma-derived human plasminogen indicated for the treatment of patients with plasminogen deficiency type 1 (hypoplasminogenemia).

PROCEDURE

A. Initial Authorization Criteria:

1. Plasminogen Deficiency Type 1 (must meet all of the following):

- Member is \geq 11 months of age
- Prescribed by or in consultation with a hematologist
- Diagnosis of hypoplasminogenemia confirmed by ALL of the following:
 - Plasminogen (PLG) gene mutation
 - Plasminogen activity levels \leq 45%
 - Documentation history of visible or non-visible lesions and symptoms consistent with plasminogen deficiency type 1
 - Abnormal wound healing OR
 - Respiratory distress/obstruction
- Member has healing of lesions/wounds suspected as a source of a recent bleeding event
- Prescriber attests that members using concomitant antiplatelet, anticoagulants or other therapies that impact normal coagulation processes will be monitored for at least 4 hours following Ryplazim infusion

B. Must be prescribed at a dose within the manufacturer's dosing guidelines (based on diagnosis, weight, etc) listed in the FDA approved labeling.

C. Ryplazim will be considered investigational or experimental for any other use and will not be covered.

D. Reauthorization Criteria:

All prior authorization renewals are reviewed on an annual basis to determine the Medical Necessity for continuation of therapy. Authorization may be extended at 1-year

intervals based upon all of the following:

- MPC Renewal:
 - Chart documentation confirming positive response to therapy as evidenced by:
 - Reduction in lesion number or size
 - Plasminogen activity trough level has increased at least 10% from baseline
 - Improvement in wound healing
 - Prescribed by or in consultation with a hematologist
 - Prescriber attests that members using concomitant antiplatelet, anticoagulants or other therapies that impact normal coagulation processes will be monitored for at least 4 hours following Ryplazim infusion

- Renewal from Previous Insurer:
 - Members who have received prior approval (from insurer other than MPC) and have been taking Ryplazim or have been receiving medication samples, should be considered under criterion A (Initial Authorization Criteria). Prescribed by or in consultation with a hematologist
 - Provider has documented positive clinical response of the member’s treatment:
 - Reduction in lesion number or size
 - Plasminogen activity trough level has increased at least 10% from baseline
 - Improvement in wound healing

Limitations:

Length of Authorization (if above criteria met)	
Initial Authorization	3 months
Reauthorization	12 months

Codes:

Code	Description
J2998	Injection, plasminogen, human-tvmh, 1mg

REFERENCES

1. Ryplazim [package insert]. Fort Lee, NJ: Prometic Bioproduction Inc; November 2021.



Ryplazim
POLICY NUMBER: RX.PA.067.MPC
REVISION DATE: 02/2026
PAGE NUMBER: 3 of 3

REVIEW HISTORY

DESCRIPTION OF REVIEW / REVISION	DATE APPROVED
Annual Review	02/2026
Annual Review	02/2025
<i>Annual policy review. Update to reauthorization criteria for non-MPC renewals</i>	02/2024
<i>Annual Review</i>	02/2023
<i>P&T Review</i>	11/2022
<i>New Policy</i>	10/2022